

## Appendix 4 NCI Common Toxicity Criteria (CTC) Version 2.0 1999

Toxicity	Grade				
	0	1	2	3	4
<b>ALLERGY/IMMUNOLOGY</b>					
Allergic reaction/hypersensitivity (including drug fever)	none	transient rash, drug fever < 38°C (<100.4°F)	urticaria, drug fever ≥ 38°C (≥100.4°F), and/or asymptomatic bronchospasm	symptomatic bronchospasm, requiring parenteral medication(s), with or without urticaria; allergy-related edema/angioedema	anaphylaxis
Note: Isolated urticaria, in the absence of other manifestations of an allergic or hypersensitivity reaction, is graded in the DERMATOLOGY/SKIN category.					
Allergic rhinitis (including sneezing, nasal stuffiness, postnasal drip)	none	mild, not requiring treatment	moderate, requiring treatment	-	-
Autoimmune reaction	none	serologic or other evidence of autoimmune reaction but patient is asymptomatic (e.g., vitiligo), all organ function is normal and no treatment is required	evidence of autoimmune reaction involving a non-essential organ or function (e.g., hypothyroidism), requiring treatment other than immunosuppressive drugs	reversible autoimmune reaction involving function of a major organ or other toxicity (e.g., transient colitis or anemia), requiring short-term immunosuppressive treatment	autoimmune reaction causing major grade 4 organ dysfunction; progressive and irreversible reaction; long-term administration of high-dose immunosuppressive therapy required
Also consider Hypothyroidism, Colitis, Hemoglobin, Hemolysis.					
Serum sickness	none	-	-	present	-
Urticaria is graded in the DERMATOLOGY/SKIN category if it occurs as an isolated symptom. If it occurs with other manifestations of allergic or hypersensitivity reaction, grade as Allergic reaction/hypersensitivity above.					
Vasculitis	none	mild, not requiring treatment	symptomatic, requiring medication	requiring steroids	ischemic changes or requiring amputation
Allergy/Immunology-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>AUDITORY/HEARING</b>					
Conductive hearing loss is graded as Middle ear/hearing in the AUDITORY/HEARING category.					
Earache is graded in the PAIN category.					
External auditory canal	normal	external otitis with erythema or dry desquamation	external otitis with moist desquamation	external otitis with discharge, mastoiditis	necrosis of the canal soft tissue or bone
Note: Changes associated with radiation to external ear (pinnae) are graded under Radiation dermatitis in the DERMATOLOGY/SKIN category.					
Inner ear/hearing	normal	hearing loss on audiometry only	tinnitus or hearing loss, not requiring hearing aid or treatment	tinnitus or hearing loss, correctable with hearing aid or treatment	severe unilateral or bilateral hearing loss (deafness), not correctable
Middle ear/hearing	normal	serous otitis without subjective decrease in hearing	serous otitis or infection requiring medical intervention; subjective decrease in hearing; rupture of tympanic membrane with discharge	otitis with discharge, mastoiditis or conductive hearing loss	necrosis of the canal soft tissue or bone
Auditory/Hearing-Other (Specify, _____)	normal	mild	moderate	severe	life-threatening or disabling
<b>BLOOD/BONE MARROW</b>					
Bone marrow cellularity	normal for age	mildly hypocellular or 25% reduction from normal cellularity for age	moderately hypocellular or >25 - ≤ 50% reduction from normal cellularity for age or >2 but <4 weeks to recovery of normal bone marrow cellularity	severely hypocellular or >50 - ≤ 75% reduction in cellularity for age or 4 - 6 weeks to recovery of normal bone marrow cellularity	aplasia or >6 weeks to recovery of normal bone marrow cellularity
Normal ranges: children (≤ 18 years) 90% cellularity average younger adults (19-59) 60-70% cellularity average older adults (≥ 60 years) 50% cellularity average					
Note: Grade Bone marrow cellularity only for changes related to treatment not disease.					
CD4 count	WNL	< LLN - 500/mm <sup>3</sup>	200 - < 500/mm <sup>3</sup>	50 - < 200/mm <sup>3</sup>	< 50/mm <sup>3</sup>
Haptoglobin	normal	decreased	-	absent	-

Toxicity	Grade				
	0	1	2	3	4
Hemoglobin (Hgb)	WNL	< LLN - 10.0 g/dl	8.0 - < 10.0 g/dl	6.5 - < 8.0 g/dl	< 6.5 g/dl
Hemolysis (e.g., immune hemolytic anemia, drug-related hemolysis, other)	none	< LLN - 100 g/L < LLN - 6.2 mmol/L only laboratory evidence of hemolysis [e.g., direct antiglobulin test (DAT, Coombs') schistocytes]	80 - < 100 g/L 4.9 - < 6.2 mmol/L evidence of red cell destruction and ≥ 2gm decrease in hemoglobin, no transfusion	65 - 80 g/L 4.0 - < 4.9 mmol/L requiring transfusion and/or medical intervention (e.g., steroids)	< 65 g/L < 4.0 mmol/L catastrophic consequences of hemolysis (e.g., renal failure, hypotension, bronchospasm, emergency splenectomy)
Also consider Haptoglobin, Hgb.					
Leukocytes (total WBC)	WNL	< LLN - $3.0 \times 10^9/L$ < LLN - 3000/mm <sup>3</sup>	≥2.0 - < $3.0 \times 10^9/L$ ≥2000 - < 3000/mm <sup>3</sup>	≥1.0 - < $2.0 \times 10^9/L$ ≥1000 - < 2000/mm <sup>3</sup>	< $1.0 \times 10^9/L$ < 1000/mm <sup>3</sup>
Lymphopenia	WNL	< LLN - $1.0 \times 10^9/L$ < LLN - 1000/mm <sup>3</sup>	≥0.5 - < $1.0 \times 10^9/L$ ≥500 - < 1000/mm <sup>3</sup>	< $0.5 \times 10^9/L$ < 500/mm <sup>3</sup>	-
Neutrophils/granulocytes (ANC/AGC)	WNL	≥1.5 - < $2.0 \times 10^9/L$ ≥1500 - < 2000/mm <sup>3</sup>	≥1.0 - < $1.5 \times 10^9/L$ ≥1000 - < 1500/mm <sup>3</sup>	≥0.5 - < $1.0 \times 10^9/L$ ≥500 - < 1000/mm <sup>3</sup>	< $0.5 \times 10^9/L$ < 500/mm <sup>3</sup>
Platelets	WNL	< LLN - < $75.0 \times 10^9/L$ < LLN - 75000/mm <sup>3</sup>	≥50.0 - < $75.0 \times 10^9/L$ ≥50000 - < 75000/mm <sup>3</sup>	≥10.0 - < $50.0 \times 10^9/L$ ≥10000 - < 50000/mm <sup>3</sup>	< $10.0 \times 10^9/L$ < 10000/mm <sup>3</sup>
Transfusion: Platelets	none	-	-	yes	platelet transfusions and other measures required to improve platelet increment; platelet transfusion refractoriness associated with life-threatening bleeding. (e.g., HLA or cross matched platelet transfusions)
Also consider Platelets.					
Transfusion: pRBCs	none	-	-	Yes	-
Also consider Hemoglobin.					
Blood/Bone Marrow-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>CARDIOVASCULAR (ARRHYTHMIA)</b>					
Conduction abnormality/ Atrioventricular heart block	none	asymptomatic, not requiring treatment (e.g., Mobitz type I second-degree AV block, Wenckebach)	symptomatic, but not requiring treatment	symptomatic and requiring treatment (e.g., Mobitz type II second-degree AV block, third-degree AV block)	life-threatening (e.g., arrhythmia associated with CHF, hypotension, syncope, shock)
Nodal/junctional arrhythmia/dysrhythmia	none	asymptomatic, not requiring treatment	symptomatic, but not requiring treatment	symptomatic and requiring treatment	life-threatening (e.g., arrhythmia associated with CHF, hypotension, syncope, shock)
Palpitations	none	present	-	-	-
Note: Grade palpitations only in the absence of a documented arrhythmia.					
Prolonged QTc interval (QTc > 0.48 seconds)	none	asymptomatic, not requiring treatment	symptomatic, but not requiring treatment	symptomatic and requiring treatment	life-threatening (e.g., arrhythmia associated with CHF, hypotension, syncope, shock)
Sinus bradycardia	none	asymptomatic, not requiring treatment	symptomatic, but not requiring treatment	symptomatic and requiring treatment	life-threatening (e.g., arrhythmia associated with CHF, hypotension, syncope, shock)
Sinus tachycardia	none	asymptomatic, not requiring treatment	symptomatic, but not requiring treatment	symptomatic and requiring treatment of underlying cause	-
Supraventricular arrhythmias (SVT/atrial fibrillation/ flutter)	none	asymptomatic, not requiring treatment	symptomatic, but not requiring treatment	symptomatic and requiring treatment	life-threatening (e.g., arrhythmia associated with CHF, hypotension, syncope, shock)
Syncope (fainting) is graded in the NEUROLOGY category.					
Vasovagal episode	none	-	present without loss of consciousness	present with loss of consciousness	-

Toxicity	Grade				
	0	1	2	3	4
Ventricular arrhythmia (PVCs/bigeminy/trigeminy/ventricular tachycardia)	none	asymptomatic, not requiring treatment	symptomatic, but not requiring treatment	symptomatic and requiring treatment	life-threatening (e.g., arrhythmia associated with CHF, hypotension, syncope, shock)
Cardiovascular/Arrhythmia-Other (Specify, _____)	none	asymptomatic, not requiring treatment	symptomatic, but not requiring treatment	symptomatic, and requiring treatment of underlying cause	life-threatening (e.g., arrhythmia associated with CHF, hypotension, syncope, shock)
<b>CARDIOVASCULAR (GENERAL)</b>					
Acute vascular leak syndrome	absent	-	symptomatic, but not requiring fluid support	respiratory compromise or requiring fluids	life-threatening; requiring pressor support and/or ventilatory support
Cardiac-ischemia/infarction	none	non-specific T-wave flattening or changes	asymptomatic, ST- and T-wave changes suggesting ischemia	angina without evidence of infarction	acute myocardial infarction
Cardiac left ventricular function	normal	asymptomatic decline of resting ejection fraction of $\geq 10\%$ but $< 20\%$ of baseline value; shortening fraction $\geq 24\%$ but $< 30\%$	asymptomatic but resting ejection fraction below LLN for laboratory or decline of resting ejection fraction $\geq 20\%$ of baseline value; $< 24\%$ shortening fraction	CHF responsive to treatment	severe or refractory CHF or requiring intubation
CNS cerebrovascular ischemia is graded in the NEUROLOGY category.					
Cardiac troponin I (cTnI)	normal	-	-	levels consistent with unstable angina as defined by the manufacturer	levels consistent with myocardial infarction as defined by the manufacturer
Cardiac troponin T (cTnT)	normal	$\geq 0.03$ - $< 0.05$ ng/ml	$\geq 0.05$ - $< 0.1$ ng/ml	$\geq 0.1$ - $< 0.2$ ng/ml	$\geq 0.2$ ng/ml
Edema	none	asymptomatic, not requiring therapy	symptomatic, requiring therapy	symptomatic edema limiting function and unresponsive to therapy or requiring drug discontinuation	anasarca (severe generalized edema)
Hypertension	none	asymptomatic, transient increase by $>20$ mmHg (diastolic) or to $> 150/100^*$ if previously WNL; not requiring treatment	recurrent or persistent or symptomatic increase by $> 20$ mmHg (diastolic) or to $> 150/100^*$ if previously WNL; not requiring treatment	requiring therapy or more intensive therapy than previously	hypertensive crisis
<i>*Note: For pediatric patients, use age and sex appropriate normal values <math>&gt; 95</math>th percentile ULN.</i>					
Hypotension	none	changes, but not requiring therapy (including transient orthostatic hypotension)	requiring brief fluid replacement or other therapy but not hospitalization; no physiologic consequences	requiring therapy and sustained medical attention, but resolves without persisting physiologic consequences	shock (associated with acidemia and impairing vital organ function due to tissue hypoperfusion)
Also consider Syncope (fainting). Note: Angina or MI is graded as Cardiac- ischemia/infarction in the CARDIOVASCULAR (GENERAL) category. <i>For pediatric patients, systolic BP 65 mmHg or less in infants up to 1 year old and 70 mmHg or less in children older than 1 year of age, use two successive or three measurements in 24 hours.</i>					
Myocarditis	none	-	-	CHF responsive to treatment	severe or refractory CHF
Operative injury of vein/artery	none	primary suture repair for injury, but not requiring transfusion	primary suture repair for injury, requiring transfusion	vascular occlusion requiring surgery or bypass for injury	myocardial infarction; resection of organ (e.g., bowel, limb)
Pericardial effusion/pericarditis	none	asymptomatic effusion, not requiring treatment	pericarditis (rub, ECG changes, and/or chest pain)	physiologic consequences resulting from symptoms	tamponade (drainage or pericardial window required)
Peripheral arterial ischemia	none	-	brief episode of ischemia managed non-surgically and without permanent deficit	requiring surgical intervention	life-threatening or with permanent functional deficit (e.g., amputation)
Phlebitis (superficial)	none	-	present	-	-
Note: Injection site reaction is graded in the DERMATOLOGY/SKIN category. Thrombosis/embolism is graded in the CARDIOVASCULAR (GENERAL) category.					
Syncope (fainting) is graded in the NEUROLOGY category.					
Thrombosis/embolism	none	-	deep vein thrombosis, not requiring anticoagulant	deep vein thrombosis, requiring anticoagulant therapy	embolic event including pulmonary embolism
Vein/artery operative injury is graded as Operative injury of vein/artery in the CARDIOVASCULAR (GENERAL) category.					

Grade					
Toxicity	0	1	2	3	4
Visceral arterial ischemia (non-myocardial)	none	-	brief episode of ischemia managed non-surgically and without permanent deficit	requiring surgical intervention	life-threatening or with permanent functional deficit (e.g., resection of ileum)
Cardiovascular/General-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>COAGULATION</b>					
Note: See the HEMORRHAGE category for grading the severity of bleeding events.					
DIC (disseminated intravascular coagulation) Also grade Platelets. Note: Must have increased fibrin split products or D-dimer in order to grade as DIC.	absent	-	-	laboratory findings present with <u>no</u> bleeding	laboratory findings <u>and</u> bleeding
Fibrinogen	WNL	≥0.75 - <1.0 x LLN	≥0.5 - <0.75 x LLN	≥0.25 - <0.5 x LLN	<0.25 x LLN
Partial thromboplastin time (PTT)	WNL	> ULN - ≤ 1.5 x ULN	> 1.5 - ≤ 2 x ULN	>2 x ULN	-
Phelbitis is graded in the CARDIOVASCULAR (GENERAL) category.					
Prothrombin time (PT)	WNL	> ULN - ≤ 1.5 x ULN	> 1.5 - ≤ 2 x ULN	>2 x ULN	-
Thrombosis/embolism is graded in the CARDIOVASCULAR (GENERAL) category.					
Thrombotic microangiopathy (e.g., thrombotic thrombocytopenic purpura/TTP or hemolytic uremic syndrome/HUS) Also consider Hemoglobin (Hgb), Platelets, Creatinine. Note: Must have microangiopathic changes on blood smear (e.g., schistocytes, helmet cells, red cell fragments).	absent	-	-	laboratory findings present without clinical consequences	laboratory findings and clinical consequences, (e.g., CNS hemorrhage/bleeding or thrombosis/embolism or renal failure) requiring therapeutic intervention
Coagulation-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>CONSTITUTIONAL SYMPTOMS</b>					
Fatigue (lethargy, malaise, asthenia) Note: See Appendix III for performance status scales.	none	increased fatigue over baseline, but not altering normal activities	moderate (e.g., decrease in performance status by 1 ECOG level or 20% Karnofsky or Lansky) <u>or</u> causing difficulty performing some activities	severe (e.g., decrease in performance status by ≥2 ECOG levels or 40% Karnofsky or Lansky) <u>or</u> loss of ability to perform some activities	bedridden or disabling
Fever (in the absence of neutropenia, where neutropenia is defined as AGC < 1.0 x 10 <sup>9</sup> /L) Also consider Allergic reaction/hypersensitivity. Note: The temperature measurements listed above are oral or tympanic.	none	38.0 - 39.0°C (100.4 - 102.2°F)	39.1 - 40.0°C (102.3 - 104.0°F)	> 40.0°C (>104.0°F) for < 24hrs	> 40.0°C (>104.0°F) for > 24hrs
Hot flashes/flushes are graded in the ENDOCRINE category.					
Rigors, chills	none	mild, requiring symptomatic treatment (e.g., blanket) or non-narcotic medication	severe and/or prolonged, requiring narcotic medication	not responsive to narcotic medication	-
Sweating (diaphoresis)	normal	mild and occasional	frequent or drenching	-	-
Weight gain Also consider Ascites, Edema, Pleural effusion.	< 5%	5 - <10%	10 - <20%	≥ 20%	-
Weight loss Also consider Vomiting, Dehydration, Diarrhea.	< 5%	5 - <10%	10 - <20%	≥20%	-
Constitutional Symptoms-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>DERMATOLOGY/SKIN</b>					
Alopecia	normal	mild hair loss	pronounced hair loss	-	-
Bruising (in absence of grade 3 or 4 thrombocytopenia) Note: Bruising resulting from grade 3 or 4 thrombocytopenia is graded as Petechiae/purpura and Hemorrhage/bleeding with grade 3 or 4 thrombocytopenia in the HEMORRHAGE category, not in the DERMATOLOGY/SKIN category.	none	localized or in dependent area	generalized	-	-

<b>Grade</b>					
<b>Toxicity</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
Dry skin	normal	controlled with emollients	not controlled with emollients	-	-
Erythema multiforme (e.g., Stevens-Johnson syndrome, toxic epidermal necrolysis)	absent	-	scattered, but not generalized eruption	severe or requiring IV fluids (e.g., generalized rash or painful stomatitis)	life-threatening (e.g., exfoliative or ulcerating dermatitis or requiring enteral or parenteral nutritional support)
Flushing	absent	present	-	-	-
Hand-foot skin reaction	none	skin changes or dermatitis without pain (e.g., erythema, peeling)	skin changes with pain, not interfering with function	skin changes with pain, interfering with function	-
Injection site reaction	none	pain or itching or erythema	pain or swelling, with inflammation or phlebitis	ulceration or necrosis that is severe or prolonged, or requiring surgery	-
Nail changes	normal	discoloration or ridging (koilonychia) or pitting	partial or complete loss of nail(s) or pain in nailbeds	-	-
Petechiae is graded in the HEMORRHAGE category.					
Photosensitivity	none	painless erythema	painful erythema	erythema with desquamation	-
Pigmentation changes (e.g., vitiligo)	none	localized pigmentation changes	generalized pigmentation changes	-	-
Pruritus	none	mild or localized, relieved spontaneously or by local measures	intense or widespread, relieved spontaneously or by systemic measures	intense or widespread and poorly controlled despite treatment	-
Purpura is graded in the HEMORRHAGE category.					
Rash/desquamation	none	macular or papular eruption or erythema without associated symptoms	macular or papular eruption or erythema with pruritus or other associated symptoms covering <50% of body surface or localized desquamation or other lesions covering <50% of body surface area	symptomatic generalized erythroderma or macular, papular or vesicular eruption or desquamation covering ≥50% of body surface area	generalized exfoliative dermatitis or ulcerative dermatitis
Also consider Allergic reaction/hypersensitivity. Note: Erythema multiforme (Stevens-Johnson syndrome) is graded separately as Erythema multiforme.					
Urticaria (hives, welts, wheals)	none	requiring no medication	requiring PO or topical treatment or IV medication or steroids for <24 hours	requiring IV medication or steroids for ≥24 hours	-
Wound- infectious	none	cellulitis	superficial infection	infection requiring IV antibiotics	necrotizing fasciitis
Wound- non-infectious	none	incisional separation	incisional hernia	fascial disruption without evisceration	fascial disruption with evisceration
Dermatology/Skin-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>ENDOCRINE</b>					
Cushingoid appearance (e.g., moon face with or without buffalo hump, centripetal obesity, cutaneous striae)	absent	-	present	-	-
Also consider Hyperglycemia, Hypokalemia.					
Feminization of male	absent	-	-	present	-
Gynecomastia	none	mild	pronounced or painful	pronounced or painful and requiring surgery	-
Hot flashes/flushes	none	mild or no more than 1 per day	moderate and greater than 1 per day	-	-
Hypothyroidism	absent	asymptomatic, TSH elevated, no therapy given	symptomatic or thyroid replacement treatment given	patient hospitalized for manifestations of hypothyroidism	myxedema coma
Masculinization of female	absent	-	-	present	-
SIADH (syndrome of inappropriate antidiuretic hormone)	absent	-	-	present	-
Endocrine-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>GASTROINTESTINAL</b>					
Amylase is graded in the METABOLIC/LABORATORY category.					

Toxicity	Grade				
	0	1	2	3	4
Anorexia	none	loss of appetite	oral intake significantly decreased	requiring IV fluids	requiring feeding tube or parenteral nutrition
Ascites (non-malignant)	none	asymptomatic	symptomatic, requiring diuretics	symptomatic, requiring therapeutic paracentesis	life-threatening physiologic consequences
Colitis	none	-	abdominal pain with mucus and/or blood in stool	abdominal pain, fever, change in bowel habits with ileus or peritoneal signs, and radiographic or biopsy documentation	perforation or requiring surgery or toxic megacolon
Also consider Hemorrhage/bleeding with grade 3 or 4 thrombocytopenia, Hemorrhage/bleeding without grade 3 or 4 thrombocytopenia, Melena/GI bleeding, Rectal bleeding/hematochezia, Hypotension.					
Constipation	none	requiring stool softener or dietary modification	requiring laxatives	obstipation requiring manual evacuation or enema	obstruction or toxic megacolon
Dehydration	none	dry mucous membranes and/or diminished skin turgor	requiring IV fluid replacement (brief)	requiring IV fluid replacement (sustained)	physiologic consequences requiring intensive care; hemodynamic collapse
Also consider Hypotension, Diarrhea, Vomiting, Stomatitis/pharyngitis (oral/pharyngeal mucositis).					
Diarrhea	none	increase of < 4 stools/day over pre-treatment	increase of 4-6 stools/day, or nocturnal stools	increase of ≥7 stools/day or incontinence; or need for parenteral support for dehydration	physiologic consequences requiring intensive care; or hemodynamic collapse
Patients without colostomy:	none				
Patients with a colostomy:	none	mild increase in loose, watery colostomy output compared with pretreatment	moderate increase in loose, watery colostomy output compared with pretreatment, but not interfering with normal activity	severe increase in loose, watery colostomy output compared with pretreatment, interfering with normal activity	physiologic consequences, requiring intensive care; or hemodynamic collapse
Duodenal ulcer (requires radiographic or endoscopic documentation)	none	-	requiring medical management or non-surgical treatment	uncontrolled by outpatient medical management; requiring hospitalization	perforation or bleeding, requiring emergency surgery
Dyspepsia/heartburn	none	mild	moderate	severe	-
Dysphagia, esophagitis, odynophagia (painful swallowing)	none	mild dysphagia, but can eat regular diet	dysphagia, requiring predominantly pureed, soft, or liquid diet	dysphagia, requiring IV hydration	complete obstruction (cannot swallow saliva) requiring enteral or parenteral nutritional support, or perforation
Note: If toxicity is radiation-related, grade either under Dysphagia- esophageal related to radiation or Dysphagia- pharyngeal related to radiation.					
Fistula- esophageal	none	-	-	present	requiring surgery
Fistula- intestinal	none	-	-	present	requiring surgery
Fistula- pharyngeal	none	-	-	present	requiring surgery
Fistula- rectal/anal	none	-	-	present	requiring surgery
Flatulence	none	mild	moderate	-	-
Gastric ulcer (requires radiographic or endoscopic documentation)	none	-	requiring medical management or non-surgical treatment	bleeding without perforation, uncontrolled by outpatient medical management; requiring hospitalization or surgery	perforation or bleeding, requiring emergency surgery
Also consider Hemorrhage/bleeding with grade 3 or 4 thrombocytopenia, Hemorrhage/bleeding without grade 3 or 4 thrombocytopenia.					
Gastritis	none	-	requiring medical management or non-surgical treatment	uncontrolled by outpatient medical management; requiring hospitalization or surgery	life-threatening bleeding, requiring emergency surgery
Also consider Hemorrhage/bleeding with grade 3 or 4 thrombocytopenia, Hemorrhage/bleeding without grade 3 or 4 thrombocytopenia.					
Hematemesis is graded in the HEMORRHAGE category.					
Hematochezia is graded in the HEMORRHAGE category as Rectal bleeding/hematochezia.					
Ileus (or neuroconstipation)	none	-	intermittent, not requiring intervention	requiring non-surgical intervention	requiring surgery
Mouth dryness	normal	mild	moderate	-	-
Mucositis					
Note: Mucositis <u>not due to radiation</u> is graded in the GASTROINTESTINAL category for specific sites: Colitis, Esophagitis, Gastritis, Stomatitis/pharyngitis (oral/pharyngeal mucositis), and Typhlitis; or the RENAL/GENITOURINARY category for Vaginitis. Radiation-related mucositis is graded as Mucositis due to radiation.					
Nausea	none	able to eat	oral intake significantly decreased	no significant intake, requiring IV fluids	-
Pancreatitis	none	-	-	abdominal pain with pancreatic enzyme elevation	complicated by shock (acute circulatory failure)
Also consider Hypotension.					
Note: Asymptomatic amylase and Amylase are graded in the METABOLIC/LABORATORY category.					
Pharyngitis is graded in the GASTROINTESTINAL category as Stomatitis/pharyngitis (oral/pharyngeal mucositis).					

Toxicity	Grade				
	0	1	2	3	4
Proctitis	none	increased stool frequency, occasional blood-streaked stools, or rectal discomfort (including hemorrhoids), not requiring medication	increased stool frequency, bleeding, mucus discharge, or rectal discomfort requiring medication; anal fissure	increased stool frequency/diarrhea, requiring parenteral support; rectal bleeding, requiring transfusion; or persistent mucus discharge, necessitating pads	perforation, bleeding or necrosis or other life-threatening complication requiring surgical intervention (e.g., colostomy)
<p>Also consider Hemorrhage/bleeding with grade 3 or 4 thrombocytopenia, Hemorrhage/bleeding without grade 3 or 4 thrombocytopenia, and Pain due to radiation.</p> <p>Note: Fistula is graded separately as Fistula- rectal/anal.</p> <p>Proctitis occurring more than 90 days after the start of radiation therapy is graded in the RTOG/EORTC Late Radiation Morbidity Scoring Scheme. (See Appendix IV)</p>					
Salivary gland changes	none	slightly thickened saliva/may have slightly altered taste (e.g., metallic); additional fluids may be required	thick, ropy, sticky saliva; markedly altered taste; alteration in diet required	-	acute salivary gland necrosis
Sense of smell	normal	slightly altered	markedly altered	-	-
Stomatitis/pharyngitis (oral/pharyngeal mucositis)	none	painless ulcers, erythema, or mild soreness in the absence of lesions	painful erythema, edema, or ulcers, but can eat or swallow	painful erythema, edema, or ulcers requiring IV hydration	severe ulceration or requires parenteral or enteral nutritional support or prophylactic intubation
<p>Note: Radiation-related mucositis is graded as Mucositis due to radiation.</p>					
Taste disturbance (dysgeusia)	normal	slightly altered	markedly altered	-	-
Typhlitis (inflammation of the cecum)	none	-	-	abdominal pain, diarrhea, fever, or radiographic documentation	perforation, bleeding or necrosis or other life-threatening complication requiring surgical intervention (e.g., colostomy)
<p>Also consider Hemorrhage/bleeding with grade 3 or 4 thrombocytopenia, Hemorrhage/bleeding without grade 3 or 4 thrombocytopenia, Hypotension, Febrile/neutropenia.</p>					
Vomiting	none	1 episode in 24 hours over pretreatment	2-5 episodes in 24 hours over pretreatment	≥6 episodes in 24 hours over pretreatment; or need for IV fluids	Requiring parenteral nutrition; or physiologic consequences requiring intensive care; hemodynamic collapse
<p>Also consider Dehydration.</p>					
<p>Weight gain is graded in the CONSTITUTIONAL SYMPTOMS category.</p>					
<p>Weight loss is graded in the CONSTITUTIONAL SYMPTOMS category.</p>					
Gastrointestinal-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>HEMORRHAGE</b>					
<p>Note: Transfusion in this section refers to pRBC infusion. For <u>any</u> bleeding with grade 3 or 4 platelets (&lt; 50,000), <u>always</u> grade Hemorrhage/bleeding with grade 3 or 4 thrombocytopenia. Also consider platelets, transfusion-pRBCs, and transfusion-platelets in addition to the grade that incorporates the site or type of bleeding. If the site or type of hemorrhage/bleeding is listed, also use the grading that incorporates the site of bleeding: CNS hemorrhage/bleeding, Hematuria, Hematemesis, Hemoptysis, Hemorrhage/bleeding with surgery, Melena/lower GI bleeding, Petechiae/purpura (Hemorrhage/bleeding into skin), Rectal bleeding/hematochezia, Vaginal bleeding. If the platelet count is ≥50,000 and the site or type of bleeding is listed, grade the specific site. If the site or type is <u>not</u> listed and the platelet count is ≥50,000, grade Hemorrhage/bleeding without grade 3 or 4 thrombocytopenia and specify the site or type in the OTHER category.</p>					
Hemorrhage/bleeding with grade 3 or 4 thrombocytopenia	none	mild without transfusion	-	requiring transfusion	catastrophic bleeding, requiring major non-elective intervention
<p>Also consider Platelets, Hemoglobin, Transfusion-platelet, Transfusion-pRBCs.</p> <p>Note: This toxicity must be graded for any bleeding with grade 3 or 4 thrombocytopenia. Also grade the site or type of hemorrhage/bleeding. If the site is not listed, grade as Other in the HEMORRHAGE category.</p>					
Hemorrhage/bleeding without grade 3 or 4 thrombocytopenia	none	mild without transfusion	-	requiring transfusion	catastrophic bleeding requiring major non-elective intervention
<p>Also consider Platelets, Hemoglobin, Transfusion-platelet, Transfusion-pRBCs.</p> <p>Note: Bleeding in the absence of grade 3 or 4 thrombocytopenia is graded here only if the specific site or type of bleeding is not listed elsewhere in the HEMORRHAGE category. Also grade as Other in the HEMORRHAGE category.</p>					
CNS hemorrhage/bleeding	none	-	-	bleeding noted on CT or other scan with no clinical consequences	hemorrhagic stroke or hemorrhagic vascular event (CVA) with neurologic signs and symptoms
Epistaxis	none	mild without transfusion	-	requiring transfusion	catastrophic bleeding, requiring major non-elective intervention
Hematemesis	none	mild without transfusion	-	requiring transfusion	catastrophic bleeding, requiring major non-elective intervention

Toxicity	Grade				
	0	1	2	3	4
Hematuria (in the absence of vaginal bleeding)	none	microscopic only	intermittent gross bleeding, no clots	persistent gross bleeding or clots; may require catheterization or instrumentation, or transfusion	open surgery or necrosis or deep bladder ulceration
Hemoptysis	none	mild without transfusion	-	requiring transfusion	catastrophic bleeding, requiring major non-elective intervention
Hemorrhage/bleeding associated with surgery Note: Expected blood loss at the time of surgery is not graded as a toxicity.	none	mild without transfusion	-	requiring transfusion	catastrophic bleeding, requiring major non-elective intervention
Melena/GI bleeding	none	mild without transfusion	-	requiring transfusion	catastrophic bleeding, requiring major non-elective intervention
Petechiae/purpura (hemorrhage/bleeding into skin or mucosa)	none	rare petechiae of skin	petechiae or purpura in dependent areas of skin	generalized petechiae or purpura of skin or petechiae of any mucosal site	-
Rectal bleeding/hematochezia	none	mild without transfusion or medication	persistent, requiring medication (e.g., steroid suppositories) and/or break from radiation treatment	requiring transfusion	catastrophic bleeding, requiring major non-elective intervention
Vaginal bleeding	none	spotting, requiring < 2 pads per day	requiring ≥ 2 pads per day, but not requiring transfusion	requiring transfusion	catastrophic bleeding, requiring major non-elective intervention
Hemorrhage-Other (Specify site, _____)	none	mild without transfusion	-	requiring transfusion	catastrophic bleeding, requiring major non-elective intervention
<b>HEPATIC</b>					
Alkaline phosphatase	WNL	> ULN - 2.5 x ULN	> 2.5 - 5.0 x ULN	> 5.0 - 20.0 x ULN	> 20.0 x ULN
Bilirubin	WNL	> ULN - 1.5 x ULN	> 1.5 - 3.0 x ULN	> 3.0 - 10.0 x ULN	> 10.0 x ULN
GGT (γ - Glutamyl transpeptidase)	WNL	> ULN - 2.5 x ULN	> 2.5 - 5.0 x ULN	> 5.0 - 20.0 x ULN	> 20.0 x ULN
Hepatic enlargement Note: Grade Hepatic enlargement only for changes related to VOD or other treatment related toxicity.	absent	-	-	present	-
Hypoalbuminemia	WNL	<LLN - 3 g/dl	≥2 - <3 g/dl	<2 g/dl	-
Liver dysfunction/failure (clinical) Note: Documented viral hepatitis is graded in the INFECTION category.	normal	-	-	asterixis	encephalopathy or coma
Portal vein flow	normal	-	decreased portal vein flow	reversal/retrograde portal vein flow	-
SGOT (AST) (serum glutamic oxaloacetic transaminase)	WNL	> ULN - 2.5 x ULN	> 2.5 - 5.0 x ULN	> 5.0 - 20.0 x ULN	> 20.0 x ULN
SGPT (ALT) (serum glutamic pyruvic transaminase)	WNL	> ULN - 2.5 x ULN	> 2.5 - 5.0 x ULN	> 5.0 - 20.0 x ULN	> 20.0 x ULN
Hepatic-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>INFECTION/FEBRILE NEUTROPENIA</b>					
Catheter-related infection	none	mild, no active treatment	moderate, localized infection, requiring local or oral treatment	severe, systemic infection, requiring IV antibiotic or antifungal treatment or hospitalization	life-threatening sepsis (e.g., septic shock)

Toxicity	Grade				
	0	1	2	3	4
Febrile neutropenia (fever of unknown origin without clinically or microbiologically documented infection) (ANC < 1.0 x 10 <sup>9</sup> /L, fever ≥38.5°C) Note: Hypothermia instead of fever may be associated with neutropenia and is graded here.	none	-	-	Present	Life-threatening sepsis (e.g., septic shock)
Infection (documented clinically or microbiologically) with grade 3 or 4 neutropenia (ANC < 1.0 x 10 <sup>9</sup> /L) Note: Hypothermia instead of fever may be associated with neutropenia and is graded here. In the absence of documented infection with grade 3 or 4 neutropenia, grade as Febrile neutropenia.	none	-	-	present	life-threatening sepsis (e.g., septic shock)
Infection with unknown ANC Note: This toxicity criterion is used in the rare case when ANC is unknown.	none	-	-	present	life-threatening sepsis (e.g., septic shock)
Infection without neutropenia	none	mild, no active treatment	moderate, localized infection, requiring local or oral treatment	severe, systemic infection, requiring IV antibiotic or antifungal treatment, or hospitalization	life-threatening sepsis (e.g., septic shock)
Infection/Febrile Neutropenia-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
Wound-infectious is graded in the DERMATOLOGY/SKIN category.					
<b>LYMPHATICS</b>					
Lymphatics	normal	mild lymphedema	moderate lymphedema requiring compression; lymphocyst	severe lymphedema limiting function; lymphocyst requiring surgery	severe lymphedema limiting function with ulceration
Lymphatics-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>METABOLIC/LABORATORY</b>					
Acidosis (metabolic or respiratory)	normal	pH < normal, but ≥7.3	-	pH < 7.3	pH < 7.3 with life-threatening physiologic consequences
Alkalosis (metabolic or respiratory)	normal	pH > normal, but ≤7.5	-	pH > 7.5	pH > 7.5 with life-threatening physiologic consequences
Amylase	WNL	> ULN - 1.5 x ULN	> 1.5 - 2.0 x ULN	> 2.0 - 5.0 x ULN	>5.0 x ULN
Bicarbonate	WNL	< LLN - 16 mEq/dl	11 - 15 mEq/dl	8 - 10 mEq/dl	< 8 mEq/dl
CPK (creatinine phosphokinase)	WNL	> ULN - 2.5 x ULN	> 2.5 - 5 x ULN	> 5 - 10 x ULN	> 10 x ULN
Hypercalcemia	WNL	> ULN - 11.5 mg/dl > ULN - 2.9 mmol/L	>11.5 - 12.5 mg/dl > 2.9 - 3.1 mmol/L	>12.5 - 13.5 mg/dl > 3.1 - 3.4 mmol/L	> 13.5 mg/dl > 3.4 mmol/L
Hypercholesterolemia	WNL	> ULN - 300 mg/dl > ULN - 7.75 mmol/L	> 300 - 400 mg/dl > 7.75 - 10.34 mmol/L	> 400 - 500 mg/dl >10.34 - 12.92 mmol/L	> 500 mg/dl > 12.92 mmol/L
Hyperglycemia	WNL	> ULN - 160 mg/dl > ULN - 8.9 mmol/L	> 160 - 250 mg/dl > 8.9 - 13.9 mmol/L	> 250 - 500 mg/dl > 13.9 - 27.8 mmol/L	> 500 mg/dl > 27.8 mmol/L or ketoacidosis
Hyperkalemia	WNL	> ULN - 5.5 mmol/L	> 5.5 - 6.0 mmol/L	> 6.0 - 7.0 mmol/L	> 7.0 mmol/L
Hypermagnesemia	WNL	> ULN - 3.0 mg/dl > ULN - 1.23 mmol/L	-	> 3.0 - 8.0 mg/dl > 1.23 - 3.30 mmol/L	> 8.0 mg/dl > 3.30 mmol/L
Hypernatremia	WNL	> ULN - 150 mmol/L	>150 - 155 mmol/L	>155 - 160 mmol/L	>160 mmol/L
Hypertriglyceridemia	WNL	> ULN - 2.5 x ULN	> 2.5 - 5.0 x ULN	> 5.0 - 10 x ULN	> 10 x ULN
Hyperuricemia	WNL	> ULN - ≤ 10 mg/dl ≤ 0.59 mmol/L without physiologic consequences	-	> ULN - ≤ 10 mg/dl ≤ 0.59 mmol/L with physiologic consequences	> 10 mg/dl > 0.59 mmol/L
Also consider Tumor lysis syndrome, Renal failure, Creatinine, Potassium.					
Hypocalcemia	WNL	<LLN - 8.0 mg/dl <LLN - 2.0 mmol/L	7.0 - < 8.0 mg/dl 1.75 - < 2.0 mmol/L	6.0 - < 7.0 mg/dl 1.5 - < 1.75 mmol/L	<6.0 mg/dl < 1.5 mmol/L
Hypoglycemia	WNL	<LLN - 55 mg/dl <LLN - 3.0 mmol/L	40 - < 55 mg/dl 2.2 - < 3.0 mmol/L	30 - < 40 mg/dl 1.7 - < 2.2 mmol/L	< 30 mg/dl < 1.7 mmol/L
Hypokalemia	WNL	<LLN - 3.0 mmol/L	-	2.5 - <3.0 mmol/L	<2.5 mmol/L
Hypomagnesemia	WNL	<LLN - 1.2 mg/dl <LLN - 0.5 mmol/L	0.9 - <1.2 mg/dl 0.4 - < 0.5 mmol/L	0.7 - < 0.9 mg/dl 0.3 - < 0.4 mmol/L	< 0.7 mg/dl < 0.3 mmol/L

<b>Grade</b>					
<b>Toxicity</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
Hyponatremia	WNL	<LLN - 130 mmol/L	-	120 - <130 mmol/L	<120 mmol/L
Hypophosphatemia	WNL	<LLN -2.5 mg/dl <LLN - 0.8 mmol/L	≥2.0 - <2.5 mg/dl ≥0.6 - <0.8 mmol/L	≥1.0 - <2.0 mg/dl ≥0.3 - <0.6 mmol/L	< 1.0 mg/dl <0.3 mmol/L
Hypothyroidism is graded in the ENDOCRINE category.					
Lipase	WNL	> ULN - 1.5 x ULN	> 1.5 - 2.0 x ULN	> 2.0 - 5.0 x ULN	> 5.0 x ULN
Metabolic/Laboratory-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>MUSCULOSKELETAL</b>					
Arthralgia is graded in the PAIN category.					
Arthritis	none	mild pain with inflammation, erythema or joint swelling but not interfering with function	moderate pain with inflammation, erythema, or joint swelling interfering with function, but not interfering with activities of daily living	severe pain with inflammation, erythema, or joint swelling and interfering with activities of daily living	disabling
Muscle weakness (not due to neuropathy)	normal	asymptomatic with weakness on physical exam	symptomatic and interfering with function, but not interfering with activities of daily living	symptomatic and interfering with activities of daily living	bedridden or disabling
Myalgia is graded in the PAIN category.					
Myositis (inflammation/damage of muscle)	none	mild pain, not interfering with function	pain interfering with function, but not interfering with activities of daily living	pain interfering with function and interfering with activities of daily living	bedridden or disabling
Also consider CPK. Note: Myositis implies muscle damage (i.e., elevated CPK).					
Osteonecrosis (avascular necrosis)	none	asymptomatic and detected by imaging only	symptomatic and interfering with function, but not interfering with activities of daily living	symptomatic and interfering with activities of daily living	symptomatic; or disabling
Musculoskeletal-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>NEUROLOGY</b>					
Aphasia, receptive and/or expressive, is graded under Speech impairment in the NEUROLOGY category.					
Arachnoiditis/meningismus/radiculitis	absent	mild pain not interfering with function	moderate pain interfering with function, but not interfering with activities of daily living	severe pain interfering with activities of daily living	unable to function or perform activities of daily living; bedridden; paraplegia
Also consider Headache, Vomiting, Fever.					
Ataxia (incoordination)	normal	asymptomatic but abnormal on physical exam, and not interfering with function	mild symptoms interfering with function, but not interfering with activities of daily living	moderate symptoms interfering with activities of daily living	bedridden or disabling
CNS cerebrovascular ischemia	none	-	-	transient ischemic event or attack (TIA)	permanent event (e.g., cerebral vascular accident)
CNS hemorrhage/bleeding is graded in the HEMORRHAGE category.					
Confusion	normal	confusion or disorientation or attention deficit of brief duration; resolves spontaneously with no sequelae	confusion or disorientation or attention deficit interfering with function, but not interfering with activities of daily living	confusion or delirium interfering with activities of daily living	harmful to others or self; requiring hospitalization
Cranial neuropathy is graded in the NEUROLOGY category as Neuropathy-cranial.					
Delusions	normal	-	-	present	toxic psychosis
Depressed level of consciousness	normal	somnolence or sedation not interfering with function	somnolence or sedation interfering with function, but not interfering with activities of daily living	obtundation or stupor; difficult to arouse; interfering with activities of daily living	coma
Note: Syncope (fainting) is graded in the NEUROLOGY category.					
Dizziness/lightheadedness	none	not interfering with function	interfering with function, but not interfering with activities of daily living	interfering with activities of daily living	bedridden or disabling
Dysphasia, receptive and/or expressive, is graded under Speech impairment in the NEUROLOGY category.					
Extrapyramidal/involuntary movement/restlessness	none	mild involuntary movements not interfering with function	moderate involuntary movements interfering with function, but not interfering with activities of daily living	severe involuntary movements or torticollis interfering with activities of daily living	bedridden or disabling
Hallucinations	normal	-	-	present	toxic psychosis
Headache is graded in the PAIN category.					
Insomnia	normal	occasional difficulty sleeping not interfering with function	difficulty sleeping interfering with function, but not interfering with activities of daily living	frequent difficulty sleeping, interfering with activities of daily living	-

Toxicity	Grade				
	0	1	2	3	4
Note: This toxicity is graded when insomnia is related to treatment. If pain or other symptoms interfere with sleep do NOT grade as insomnia.					
Memory loss	normal	memory loss not interfering with function	memory loss interfering with function, but not interfering with activities of daily living	memory loss interfering with activities of daily living	amnesia
Mood alteration-anxiety agitation	normal	mild mood alteration not interfering with function	moderate mood alteration interfering with function, but not interfering with activities of daily living	severe mood alteration interfering with activities of daily living	suicidal ideation or danger to self
Mood alteration-depression	normal	mild mood alteration not interfering with function	moderate mood alteration interfering with function, but not interfering with activities of daily living	severe mood alteration interfering with activities of daily living	suicidal ideation or danger to self
Mood alteration-euphoria	normal	mild mood alteration not interfering with function	moderate mood alteration interfering with function, but not interfering with activities of daily living	severe mood alteration interfering with activities of daily living	danger to self
Neuropathic pain is graded in the PAIN category.					
Neuropathy- cranial	absent	-	present, not interfering with activities of daily living	present, interfering with activities of daily living	life-threatening, disabling
Neuropathy- motor	normal	subjective weakness but no objective findings	mild objective weakness interfering with function, but not interfering with activities of daily living	objective weakness interfering with activities of daily living	paralysis
Neuropathy-sensory	normal	loss of deep tendon reflexes or paresthesia (including tingling) but not interfering with function	objective sensory loss or paresthesia (including tingling), interfering with function, but not interfering with activities of daily living	sensory loss or paresthesia interfering with activities of daily living	permanent sensory loss that interferes with function
Nystagmus Also consider Vision-double vision.	absent	present	-	-	-
Personality/behavioral	normal	change, but not disruptive to patient or family	disruptive to patient or family	disruptive to patient and family; requiring mental health intervention	harmful to others or self; requiring hospitalization
Pyramidal tract dysfunction (e.g., ↑ tone, hyperreflexia, positive Babinski, ↓ fine motor coordination)	normal	asymptomatic with abnormality on physical examination	symptomatic or interfering with function but not interfering with activities of daily living	interfering with activities of daily living	bedridden or disabling; paralysis
Seizure(s)	none	-	seizure(s) self-limited and consciousness is preserved	seizure(s) in which consciousness is altered	seizures of any type which are prolonged, repetitive, or difficult to control (e.g., status epilepticus, intractable epilepsy)
Speech impairment (e.g., dysphasia or aphasia)	normal	-	awareness of receptive or expressive dysphasia, not impairing ability to communicate	receptive or expressive dysphasia, impairing ability to communicate	inability to communicate
Syncope (fainting) Also consider CARDIOVASCULAR (ARRHYTHMIA), Vasovagal episode, CNS cerebrovascular ischemia.	absent	-	-	present	-
Tremor	none	mild and brief or intermittent but not interfering with function	moderate tremor interfering with function, but not interfering with activities of daily living	severe tremor interfering with activities of daily living	-
Vertigo	none	not interfering with function	interfering with function, but not interfering with activities of daily living	interfering with activities of daily living	bedridden or disabling
Neurology-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>OCULAR/VISUAL</b>					
Cataract	none	asymptomatic	symptomatic, partial visual loss	symptomatic, visual loss requiring treatment or interfering with function	-
Conjunctivitis	none	abnormal ophthalmologic changes, but asymptomatic or symptomatic without visual impairment (i.e., pain and irritation)	symptomatic and interfering with function, but not interfering with activities of daily living	symptomatic and interfering with activities of daily living	-
Dry eye	normal	mild, not requiring treatment	moderate or requiring artificial tears	-	-
Glaucoma	none	increase in intraocular pressure but no visual loss	increase in intraocular pressure with retinal changes	visual impairment	unilateral or bilateral loss of vision (blindness)

<b>Grade</b>					
<b>Toxicity</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
Keratitis (corneal inflammation/corneal ulceration)	none	abnormal ophthalmologic changes but asymptomatic or symptomatic without visual impairment (i.e., pain and irritation)	symptomatic and interfering with function, but not interfering with activities of daily living	symptomatic and interfering with activities of daily living	unilateral or bilateral loss of vision (blindness)
Tearing (watery eyes)	none	mild: not interfering with function	moderate: interfering with function, but not interfering with activities of daily living	interfering with activities of daily living	-
Vision- blurred vision	normal	-	symptomatic and interfering with function, but not interfering with activities of daily living	symptomatic and interfering with activities of daily living	-
Vision- double vision (diplopia)	normal	-	symptomatic and interfering with function, but not interfering with activities of daily living	symptomatic and interfering with activities of daily living	-
Vision- flashing lights/floaters	normal	mild, not interfering with function	symptomatic and interfering with function, but not interfering with activities of daily living	symptomatic and interfering with activities of daily living	-
Vision- night blindness (nyctalopia)	normal	abnormal electroretinography but asymptomatic	symptomatic and interfering with function, but not interfering with activities of daily living	symptomatic and interfering with activities of daily living	-
Vision- photophobia	normal	-	symptomatic and interfering with function, but not interfering with activities of daily living	symptomatic and interfering with activities of daily living	-
Ocular/Visual-Other (Specify, _____)	normal	mild	moderate	severe	unilateral or bilateral loss of vision (blindness)
<b>PAIN</b>					
Abdominal pain or cramping	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Arthralgia (joint pain)	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Arthritis (joint pain with clinical signs of inflammation) is graded in the MUSCULOSKELETAL category.					
Bone pain	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Chest pain (non-cardiac and non-pleuritic)	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Dysmenorrhea	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Dyspareunia	none	mild pain not interfering with function	moderate pain interfering with sexual activity	severe pain preventing sexual activity	-
Dysuria is graded in the RENAL/GENITOURINARY category.					
Earache (otalgia)	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Headache	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Hepatic pain	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Myalgia (muscle pain)	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling

Toxicity	Grade				
	0	1	2	3	4
Neuropathic pain (e.g., jaw pain, neurologic pain, phantom limb pain, post-infectious neuralgia, or painful neuropathies)	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Pelvic pain	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Pleuritic pain	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Rectal or perirectal pain (proctalgia)	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Tumor pain (onset or exacerbation of tumor pain due to treatment)	none	mild pain not interfering with function	moderate pain: pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain: pain or analgesics severely interfering with activities of daily living	disabling
Tumor flair is graded in the SYNDROME category.					
Pain-Other (Specify, _____)	none	mild	moderate	severe	disabling
<b>PULMONARY</b>					
Adult Respiratory Distress Syndrome (ARDS)	absent	-	-	-	present
Apnea	none	-	-	present	requiring intubation
Carbon monoxide diffusion capacity (DL <sub>CO</sub> )	≥ 90% of pretreatment or normal value	≥75 - <90% of pretreatment or normal value	≥50 - <75% of pretreatment or normal value	≥25 - <50% of pretreatment or normal value	< 25% of pretreatment or normal value
Cough	absent	mild, relieved by non-prescription medication	requiring narcotic antitussive	severe cough or coughing spasms, poorly controlled or unresponsive to treatment	-
Dyspnea (shortness of breath)	normal	-	dyspnea on exertion	dyspnea at normal level of activity	dyspnea at rest or requiring ventilator support
FEV <sub>1</sub>	≥ 90% of pretreatment or normal value	≥75 - <90% of pretreatment or normal value	≥50 - <75% of pretreatment or normal value	≥25 - <50% of pretreatment or normal value	< 25% of pretreatment or normal value
Hiccoughs (hiccups, singultus)	none	mild, not requiring treatment	moderate, requiring treatment	severe, prolonged, and refractory to treatment	-
Hypoxia	normal	-	decreased O <sub>2</sub> saturation with exercise	decreased O <sub>2</sub> saturation at rest, requiring supplemental oxygen	decreased O <sub>2</sub> saturation, requiring pressure support (CPAP) or assisted ventilation
Pleural effusion (non-malignant)	none	asymptomatic and not requiring treatment	symptomatic, requiring diuretics	symptomatic, requiring O <sub>2</sub> or therapeutic thoracentesis	life-threatening (e.g., requiring intubation)
Pleuritic pain is graded in the PAIN category.					
Pneumonitis/pulmonary infiltrates	none	radiographic changes but asymptomatic or symptoms not requiring steroids	radiographic changes and requiring steroids or diuretics	radiographic changes and requiring oxygen	radiographic changes and requiring assisted ventilation
Pneumothorax	none	no intervention required	chest tube required	sclerosis or surgery required	life-threatening
Pulmonary embolism is graded as Thrombosis/embolism in the CARDIOVASCULAR (GENERAL) category.					
Pulmonary fibrosis	none	radiographic changes, but asymptomatic or symptoms not requiring steroids	requiring steroids or diuretics	requiring oxygen	requiring assisted ventilation
Note: Radiation-related pulmonary fibrosis is graded in the RTOG/EORTC Late Radiation Morbidity Scoring Scheme- Lung. (See Appendix IV)					
Voice changes/stridor/larynx (e.g., hoarseness, loss of voice, laryngitis)	normal	mild or intermittent hoarseness	persistent hoarseness, but able to vocalize; may have mild to moderate edema	whispered speech, not able to vocalize; may have marked edema	marked dyspnea/stridor requiring tracheostomy or intubation
Note: Cough from radiation is graded as cough in the PULMONARY category. Radiation-related hemoptysis from larynx/pharynx is graded as Grade 4 Mucositis due to radiation in the GASTROINTESTINAL category. Radiation-related hemoptysis from the thoracic cavity is graded as Grade 4 Hemoptysis in the HEMORRHAGE category.					

Toxicity	Grade				
	0	1	2	3	4
Pulmonary-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>RENAL/GENITOURINARY</b>					
Bladder spasms	absent	mild symptoms, not requiring intervention	symptoms requiring antispasmodic	severe symptoms requiring narcotic	-
Creatinine	WNL	> ULN - 1.5 x ULN	> 1.5 - 3.0 x ULN	> 3.0 - 6.0 x ULN	> 6.0 x ULN
<i>Note: Adjust to age-appropriate levels for pediatric patients.</i>					
Dysuria (painful urination)	none	mild symptoms requiring no intervention	symptoms relieved with therapy	symptoms not relieved despite therapy	-
Fistula or GU fistula (e.g., vaginal, vesicovaginal)	none	-	-	requiring intervention	requiring surgery
Hemoglobinuria	-	present	-	-	-
Hematuria (in the absence of vaginal bleeding) is graded in the HEMORRHAGE category.					
Incontinence	none	with coughing, sneezing, etc.	spontaneous, some control	no control (in the absence of fistula)	-
Operative injury to bladder and/or ureter	none	-	injury of bladder with primary repair	sepsis, fistula, or obstruction requiring secondary surgery; loss of one kidney; injury requiring anastomosis or re-implantation	septic obstruction of both kidneys or vesicovaginal fistula requiring diversion
Proteinuria	normal or < 0.15 g/24 hours	1+ or 0.15 - 1.0 g/24 hours	2+ to 3+ or 1.0 - 3.5 g/24 hours	4+ or > 3.5 g/24 hours	nephrotic syndrome
<i>Note: If there is an inconsistency between absolute value and uristix reading, use the absolute value for grading.</i>					
Renal failure	none	-	-	requiring dialysis, but reversible	requiring dialysis and irreversible
Ureteral obstruction	none	unilateral, not requiring surgery	-	bilateral, not requiring surgery	stent, nephrostomy tube, or surgery
Urinary electrolyte wasting (e.g., Fanconi's syndrome, renal tubular acidosis)	none	asymptomatic, not requiring treatment	mild, reversible and manageable with oral replacement	reversible but requiring IV replacement	irreversible, requiring continued replacement
<i>Also consider Acidosis, Bicarbonate, Hypocalcemia, Hypophosphatemia.</i>					
Urinary frequency/urgency	normal	increase in frequency or nocturia up to 2 x normal	increase > 2 x normal but < hourly	hourly or more with urgency, or requiring catheter	-
Urinary retention	normal	hesitancy or dribbling, but no significant residual urine; retention occurring during the immediate postoperative period	hesitancy requiring medication or occasional in/out catheterization (<4 x per week), or operative bladder atony requiring indwelling catheter beyond immediate postoperative period but for < 6 weeks	requiring frequent in/out catheterization (≥ 4 x per week) or urological intervention (e.g., TURP, suprapubic tube, urethrotomy)	bladder rupture
Urine color change (not related to other dietary or physiologic cause e.g., bilirubin, concentrated urine, hematuria)	normal	asymptomatic, change in urine color	-	-	-
Vaginal bleeding is graded in the HEMORRHAGE category.					
Vaginitis (not due to infection)	none	mild, not requiring treatment	moderate, relieved with treatment	severe, not relieved with treatment, or ulceration not requiring surgery	ulceration requiring surgery
Renal/Genitourinary -Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling
<b>SECONDARY MALIGNANCY</b>					
Secondary Malignancy-Other (Specify type, _____) excludes metastatic tumors	none	-	-	-	present
<b>SEXUAL/REPRODUCTIVE FUNCTION</b>					
Dyspareunia is graded in the PAIN category.					
Dysmenorrhea is graded in the PAIN category.					
Erectile impotence	normal	mild (erections impaired but satisfactory)	moderate (erections impaired, unsatisfactory for intercourse)	no erections	-
Female sterility	normal	-	-	sterile	-

Toxicity	Grade				
	0	1	2	3	4
Feminization of male is graded in the ENDOCRINE category.					
Irregular menses (change from baseline)	normal	occasionally irregular or lengthened interval, but continuing menstrual cycles	very irregular, but continuing menstrual cycles	persistent amenorrhea	-
Libido	normal	decrease in interest	severe loss of interest	-	-
Male infertility	-	-	Oligospermia (low sperm count)	Azoospermia (no sperm)	-
Masculinization of female is graded in the ENDOCRINE category.					
Vaginal dryness	normal	mild	requiring treatment and/or interfering with sexual function, dyspareunia	-	-
Sexual/Reproductive Function-Other (Specify, _____)	none	mild	moderate	severe	disabling
<b>SYNDROMES (not included in previous categories)</b>					
Acute vascular leak syndrome is graded in the CARDIOVASCULAR (GENERAL) category.					
ARDS (Adult Respiratory Distress Syndrome) is graded in the PULMONARY category.					
Autoimmune reactions are graded in the ALLERGY/IMMUNOLOGY category.					
DIC (disseminated intravascular coagulation) is graded in the COAGULATION category.					
Fanconi's syndrome is graded as Urinary electrolyte wasting in the RENAL/GENITOURINARY category.					
Renal tubular acidosis is graded as Urinary electrolyte wasting in the RENAL/GENITOURINARY category.					
Stevens-Johnson syndrome (erythema multiforme) is graded in the DERMATOLOGY/SKIN category.					
SIADH (syndrome of inappropriate antidiuretic hormone) is graded in the ENDOCRINE category.					
Thrombotic microangiopathy (e.g., thrombotic thrombocytopenic purpura/TTP or hemolytic uremic syndrome/HUS) is graded in the COAGULATION category.					
Tumor flare	none	mild pain not interfering with function	moderate pain; pain or analgesics interfering with function, but not interfering with activities of daily living	severe pain; pain or analgesics interfering with function and interfering with activities of daily living	Disabling
Also consider Hypercalcemia. Note: Tumor flare is characterized by a constellation of symptoms and signs in direct relation to initiation of therapy (e.g., anti-estrogens/androgens or additional hormones). The symptoms/signs include tumor pain, inflammation of visible tumor, hypercalcemia, diffuse bone pain, and other electrolyte disturbances.					
Tumor lysis syndrome	absent	-	-	present	-
Also consider Hyperkalemia, Creatinine.					
Urinary electrolyte wasting (e.g., Fanconi's syndrome, renal tubular acidosis) is graded under the RENAL/GENITOURINARY category.					
Syndromes-Other (Specify, _____)	none	mild	moderate	severe	life-threatening or disabling